## National Climatic Data Center

## DATA DOCUMENTATION

## FOR

# **DATA SET 6141A (DSI-6141A)**

Eta Data Assimilation System (EDAS)
Archive Information

December 6, 2002

National Climatic Data Center 151 Patton Ave. Asheville, NC 28801-5001 USA

### Table of Contents

Top	ic	Page	Numb∈	٤r
1.	Abstract			3
2.	Element Names and Definitions:			4
3.	Start Date			6
4.	Stop Date			6
5.	Coverage			6
6.	How to order data			6
7.	Archiving Data Center			7
8.	Technical Contact			7
9.	Known Uncorrected Problems			7
10.	Quality Statement			7
11.	Essential Companion Data Sets			7
12	References			7

1. Abstract: The National Weather Service's National Centers for Environmental Prediction (NCEP) runs a series of computer analyses and forecasts operationally. One of the operational systems is the Global Data Assimilation System (GDAS, Kanamitsu, 1989), which uses the spectral Medium Range Forecast model (MRF) for the forecast. Another system is the Eta Data Assimilation System (EDAS), covering the U.S.

At NOAA's <u>Air Resources Laboratory</u> (ARL), NCEP model output are used for air quality transport and dispersion modeling. ARL archives both EDAS and GDAS data using a 1-byte packing method. Both archives contain basic fields such as the u- and v-wind components, temperature, and humidity. However, the archives differ from each other because of the horizontal and vertical resolution, as well as in the specific fields, provided by NCEP.

The archive data file contains the data in synoptic time sequence, without any missing records (missing data is represented by nulls and the forecast hour is set to negative 1). Therefore it is possible to position randomly to any point within a data file. Each file contains data for approximately two weeks: days one through 15, and 16 through the end of the month. At each time period, an index record is always the first record, followed by surface data, and then all data in each pressure level from the ground up.

#### Data Grid

The data are on a 79 by 55 Lambert Conformal grid (Fig. 1). In Table 1, the data grid is identified by the model that produced the data, a grid identification number, the number of X and Y grid points, the Pole position (latitude and longitude) of the grid projection, a reference latitude and longitude, the grid spacing (km) which is true at the reference point, the orientation with respect to the reference longitude, the angle between the axis and the cone, and a point on the grid in grid units and latitude and longitude. The given pole position results in the lowest left grid point to have a value of (1,1).

		 		Table	= 1. Dat	a Grid	d Speci	ficat	ions					
					Pole Lon.						-	-	_	-
EDAS	38	 79	55	90N	0W	35N	95W	80	0	25	39.0	25.0	35N	95W

#### ORIGIN OF THE DATA

The 3-hourly archive data come from NCEP's EDAS. The EDAS was implemented into the operational early Eta model runs during 1995. The EDAS is an intermittent assimilation system consisting of successive 3-h Eta model forecasts and Optimum Interpolation (OI) analyses for a pre-forecast period (12-h for the early Eta) on a 38 level, 48 km grid. A 6-h forecast from the GDAS is used to to start the assimilation at 12-h prior to model start time. The following is a schematic for the 12Z cycle

The 3-h analysis updates allow for the use of high frequency observations, such as wind profiler, NEXRAD, and aircraft data. ARL saves the successive 3-hour analyses, twice each day to produce a continuous data archive. Some fields such as precipitation and surface fluxes are not available in the analysis files, therefore these are taken from the successive 3-hour forecast files. The 48 km data are interpolated to a 40 km, Lambert Conformal Grid, covering the continental United States.

#### ARL PROCESSING

The ARL archiving program extracts every other grid point of the 3 hourly, 40 km data to produce a 3 hourly, 80 km dataset on pressure surfaces. In addition, 14 gridpoints on the western end of the model domain and 10 gridpoints on the northern end of the domain are removed to reduce the size of the semi-monthly files (currently at about 84 Mbytes). The data are put into semi-monthly files and saved for shipment to NCDC. About 6 months of EDAS data are also put online at ARL's web site for easy access via ftp:

http://www.arl.noaa.gov/ss/transport/archives.html

#### Element Names and Definitions:

Field	Units :	Label	Data Orde
Pressure reduced to mean sea level	hPa	MSLP	
Temperature at surface	K	TMPS	S2
Accumulated precipitation (3 h accumulation)	m	TPP3	S3
Accumulated convective precipitation (3 h accumulation)	m	CPP3	S4
Volumetric soil moisture content	frac.	SOLW	S5
Temperature at 2m AGL	K	T02M	S6
Relative Humidity at 2m AGL	%	RH2M	s7
U-component of wind at 10 m AGL	m/s	U10M	S8
V-component of wind at 10 m AGL	m/s	V10M	S9
Pressure at surface	hPa	PRSS	S10
Categorial snow (yes=1, no=0)	-	CSNO	S11
Categorial rain (yes=1, no=0)	-	CRAI	S12
Latent heat net flux at surface	W/m2	LHTF	S13
Sensible heat net flux at surface	W/m2	SHTF	S14
Momentum flux, u-component at surface	N/m2	UMOF	S15
Momentum flux, v-component at surface	N/m2	VMOF	S16
Low cloud cover	%	LCLD	S17
Medium cloud cover	%	MCLD	S18
High cloud cover	8	HCLD	S19
Total cloud cover	%	TCLD	S20
Downward short wave radiation flux	W/m2	DSWF	S21
U-component of wind with respect to grid	m/s	UWND	U1
V-component of wind with respect to grid	m/s	VWND	U2

Geopotential height	gpm*	HGTS	U3
Temperature	K	TEMP	U4
Pressure vertical velocity	hPa/s	WWND	U5
Relative humidity	용	RELH	U6

\_\_\_\_\_\_\_

#### Meteorological Fields and Vertical Structure

The archived data files contain only some of the fields normally produced by the model at NCEP. These were selected according to what is most relevant for transport and dispersion studies and disk space limitations. In Table 2, the fields are identified by a description, the units, and a unique four character identification label that is written to the header label of each record. Data order in the file is given by a two digit code. The first digit indicates if it is a surface (or single) level variable (S) or an upper level variable (U). The second digit indicates the order in which that variable appears in the file. The upper level EDAS data are output on the following 22 pressure surfaces. Table 3 gives the level number corresponding to each data level, which is also written to each header label.

Table 3. Description of Vertical Levels

Table	3. De:	scri	ption	of	Vertica	l Levels	5		
 	Level	Н (	eight					 	
 	 22	 50	hPa					 	
	21								
		150							
		200							
		250							
	17	300							
		400							
		500							
	14		hPa						
	13	600							
		650							
	11	700							
	10	750	hPa						
	9	800	hPa						
	8	825	hPa						
	7	850	hPa						
	6	875	hPa						
	5	900	hPa						
	4	925	hPa						
		950	hPa						
		975	hPa						
	1		) hPa						
	0	sur	face						

Definition File - EDAS.CFG

MODEI	L TYPE:	EDAS
GRID	NUMB:	38
VERT	COORD:	2
POLE	LAT:	90.

.

<sup>\*</sup> geopotential meters

```
POLE LON:
                        35.
REF LAT:
REF LON:
                       -95.
REF GRID:
                       80.0
ORIENTATION:
                        0.
CONE ANGLE:
                       25.
SYNC X:
SYNC Y:
                       25.0
SYNC LAT:
                       35.
SYNC LON:
                      -95.
                        0.
SPECIAL:
NUMB X:
                        79
NUMB Y:
                        55
NUMB LEVELS:
                        23
                                  21 MSLP TMPS TPP3 CPP3 SOLW T02M RH2M
LEVEL 1:
                        0.
                                      U10M V10M PRSS CSNO CRAI LHTF SHTF
                                      UMOF VMOF LCLD MCLD HCLD TCLD DSWF
                     1000.
                                 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 2:
                      975.
LEVEL 3:
                                06 UWND VWND HGTS TEMP WWND RELH
LEVEL 4:
                                06 UWND VWND HGTS TEMP WWND RELH
                      950.
                    925. 06 UWND VWND HGTS TEMP WWND RELH
900. 06 UWND VWND HGTS TEMP WWND RELH
875. 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 5:
LEVEL 6:
LEVEL 7:
                    875. 06 UWND VWND HGTS TEMP WWND RELH
850. 06 UWND VWND HGTS TEMP WWND RELH
825. 06 UWND VWND HGTS TEMP WWND RELH
800. 06 UWND VWND HGTS TEMP WWND RELH
750. 06 UWND VWND HGTS TEMP WWND RELH
700. 06 UWND VWND HGTS TEMP WWND RELH
650. 06 UWND VWND HGTS TEMP WWND RELH
600. 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 8:
LEVEL 9:
LEVEL 10:
LEVEL 11:
LEVEL 12:
LEVEL 13:
LEVEL 14:
                      550. 06 UWND VWND HGTS TEMP WWND RELH
500. 06 UWND VWND HGTS TEMP WWND RELH
400. 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 15:
LEVEL 16:
LEVEL 17:
                      300. 06 UWND VWND HGTS TEMP WWND RELH
250. 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 18:
LEVEL 19:
                      200.
LEVEL 20:
                                06 UWND VWND HGTS TEMP WWND RELH
                      150.
LEVEL 21:
                                 06 UWND VWND HGTS TEMP WWND RELH
                      100. 06 UWND VWND HGTS TEMP WWND RELH 50. 06 UWND VWND HGTS TEMP WWND RELH
LEVEL 22:
LEVEL 23:
```

3. <u>Start Date</u>: 19970101

4. Stop Date: Ongoing.

5. Coverage: North America

a. Southernmost Latitude: 25N
b. Northernmost Latitude: 50N
c. Westernmost Longitude: 125W
d. Easternmost Longitude: 65W

#### 6. How to Order Data:

Ask NCDC's Climate Services about the cost of obtaining this data set.

Phone: 828-271-4800 FAX: 828-271-4876

•

E-mail: NCDC.Orders@noaa.gov

### 7. Archiving Data Center:

National Climatic Data Center Federal Building 151 Patton Avenue Asheville, NC 28801-5001 Phone: (828) 271-4800.

#### 8. <u>Technical Contact</u>:

National Climatic Data Center Federal Building 151 Patton Avenue Asheville, NC 28801-5001 Phone: (828) 271-4800.

NOAA-Air Resources Laboratory 1315 East-West Highway Silver Spring, MD 20910 Phone: (301 713-0295)

- 9. Known Uncorrected Problems: None.
- 10. Quality Statement: No information provided with original documentation.
- 11. Essential Companion Datasets: None.

#### 12. References:

Kanamitsu, M., 1989: Description of the NMC Global Data Assimilation and Forecast System, Weather and Forecasting, 4 (335-342).

Petersen, R.A. and J.D. Stackpole, 1989: Overview of the NMC Production Suite, Weather and Forecasting, 4 (313-322).

Sela, J.G., 1980: Spectral modeling at the National Meteorological Center, Mon. Wea. Rev.,  $108 \ (1279-1292)$ .

: